



BOXER

% 80 EC

**Definite control of annual
weeds**



Active ingredient :-

Prosulfocarb 80% EC

Introduction :-

A selective weed killer, used before germination to control some slender and broadleaf weeds in wheat and barley.

Advantages of using a boxer:

It control many types of weeds. There is no restriction on the use of the sown area for the next season, that is, it does not affect crop rotation

Usage recommendations :-

- 1) Surface planting of seeds should be avoided so that the planting depth is not less than 2.5-3 cm for sandy soil and 2 cm for clay soil.**
- 2) Boxer is used after planting and before germination. After the first or second irrigation , provided that the soil has moisture, and a period of 24 hours is waited after the treatment and before starting the second irrigation .**
- 3) The greatest effectiveness of the effect is the recommended time of use, but in the later stages, the use is less effective or useless in general.**
- 4) The degree of soil moisture must be taken into account when treating, so that it does not exceed 50% of the field capacity.**
- 5) The optimal date for using Boxer is in the stage before germination and the emergence of seedlings above the soil surface, in order to avoid any damage to the wheat seedlings. for a period of 7-14 days. This is according to the surrounding weather conditions, knowing that these seedlings regain their activity later to continue their growth naturally until the end of the season.**
- 6) In the event of a high infestation of broadleaf weeds, it is recommended to use another herbicide to control broadleaf weeds at a later stage by using Boxer to ensure optimal control of these weeds.**
- 7) Like any other herbicide, it must be taken into account when using Boxer that the crop is not exposed to stress due to some external factors such as severe cold, frost, thirst, or in the event of a high percentage of nematodes in the soil.**
- 8) As is the case with other herbicides, there is the possibility of some Rottboellia weed plants appearing after using the herbicide Boxer, at a late stage of growth and development of wheat plants, knowing that these herbs do not have any economic effect on the crop, as they have no effect on the development and growth of wheat plants, because Late germination and inability to compete with the crop**

Weed examples



Vicia sativa



Sonchus oleraceus



Rumex dentatus



Chenopodium spp.



Melilotus indica



Malva parviflora



Rottboellia spp.

Usage and dosage :-

Crop	Weeds	Dose	Usage
Wheat, barley and potatoes	Thin-leaved grasses, such as henbane, are sensitive and resistant	4 L/Ha	It is used after planting and before germination, after the first or second irrigation, provided that the soil is in the degree of cultivation, and it is waited for 24 hours after treatment and before starting the next irrigation.
	Broad-leaved herbs such as mallow, zarbikh, and cock's mane	3 - 4 L/Ha	

Usage recommendations :-

- Mixing it with the tank is easy and it cannot cause clogging of the equipment .
- For mixing operations with other chemical elements, a mini experiment is recommended .
- before using it extensively .
- Avoid spraying in bad conditions such as intense sunlight - high temperatures .

Packing :-

Boxer is available in 5 liter gallons.